

*Correction*

J Geophys (1980) 48:158–160

**High Precision Measurement of the Frequency of Mode  $\circ S_o$** W. Zürn,<sup>1</sup> L. Knopoff,<sup>2</sup> and P.A. Rydelek<sup>2</sup><sup>1</sup> Geowissenschaftliches Gemeinschaftsobservatorium, Universitäten Karlsruhe/Stuttgart, Heubach 206, D-7620 Wolfach, Federal Republic of Germany<sup>2</sup> Institute of Geophysics and Planetary Physics, University of California at Los Angeles, Los Angeles, California 90024, USA

We are very grateful to Dr. Michael Shimshoni for pointing out a small numerical error (<1 s.d.) in the paper by Zürn et al. (1980). The corrected frequency of mode  $\circ S_o$  is:

$$f(\circ S_o) = 2.932827 \text{ cph} \pm 11 \text{ ppm.}$$

The new value is in agreement at the 1 s.d. level with the frequency reported by Riedsel et al. (1980).

**References**

- Riedsel, M.A., Agnew, D., Berger, J., Gilbert, F.: Stacking for the frequencies and Qs of  $\circ S_o$  and  $\circ S_o$ . Geophys. J.R. Astron. Soc. **62**, 457–471, 1980  
Zürn, W., Knopoff, L., Rydelek, P.A.: High precision measurement of the frequency of mode  $\circ S_o$ . J. Geophys. **48**, 158–160, 1980

Received April 6, 1981